

Australian Standard[®]

Sensory analysis

**Method 2.4: Specific methods—Duo-trio
test**

STANDARDS
Australia



This Australian Standard® was prepared by Committee FT-022, Sensory Analysis. It was approved on behalf of the Council of Standards Australia on 16 September 2005. This Standard was published on 12 October 2005.

The following are represented on Committee FT-022:

- Australian Associated Brewers
 - Australian Food and Grocery Council
 - Australian Institute of Food Science and Technology Limited
 - Australian Paint Manufacturers' Federation
 - Australian Society of Cosmetic Chemists
 - Australian Wine and Brandy Corporation
 - Department of Primary Industries and Fisheries Queensland
 - Food Science Australia
 - Food Technology Association of Victoria
 - National Association of Testing Authorities Australia
 - RMIT University
-

This Standard was issued in draft form for comment as DR 00333.

Standards Australia wishes to acknowledge the participation of the expert individuals that contributed to the development of this Standard through their representation on the Committee and through the public comment period.

Keeping Standards up-to-date

Australian Standards® are living documents that reflect progress in science, technology and systems. To maintain their currency, all Standards are periodically reviewed, and new editions are published. Between editions, amendments may be issued.

Standards may also be withdrawn. It is important that readers assure themselves they are using a current Standard, which should include any amendments that may have been published since the Standard was published.

Detailed information about Australian Standards, drafts, amendments and new projects can be found by visiting www.standards.org.au

Standards Australia welcomes suggestions for improvements, and encourages readers to notify us immediately of any apparent inaccuracies or ambiguities. Contact us via email at mail@standards.org.au, or write to Standards Australia, GPO Box 476, Sydney, NSW 2001.

Australian Standard[®]

Sensory analysis

Method 2.4: Specific methods—Duo-trio test

Originally as AS 2542.2.4—1988.
Second edition 2005.
Revised incorporating Amendment No. 1 (February 2008).

COPYRIGHT

© Standards Australia

All rights are reserved. No part of this work may be reproduced or copied in any form or by any means, electronic or mechanical, including photocopying, without the written permission of the publisher.

Published by Standards Australia GPO Box 476, Sydney, NSW 2001, Australia
ISBN 0 7337 6943 8

PREFACE

This Standard was prepared by the Standards Australia Committee FT-022, Sensory Analysis to supersede AS 2542.2.4—1988, *Sensory analysis of foods, Part 2.4: Specific methods—Duo-trio test*.

This Standard incorporates Amendment No. 1 (February 2008). The changes required by the Amendment are indicated in the text by a marginal bar and amendment number against the clause, note, table, figure or part thereof affected.

This Standard is identical with and reproduced from ISO 10399:2004, *Sensory analysis—Methodology—Duo-trio test*.

The objective of this Standard is to provide a procedure for determining whether a perceptible sensory difference or similarity exists between samples of two products. The method is a forced-choice procedure. The method is applicable whether a difference exists in a single sensory attribute or in several attributes.

A1 In reference to Table A.1 the exact p level for α can be calculated using binomial statistics. For example, using Microsoft Excel the p value for $\alpha = 1 - \text{BINOMDIST}(x, n, 1/2, \text{true})$ for x correct responses from n panellists.

In reference to Table A.2, the exact p level for β can be calculated using binomial statistics. For example, using Microsoft Excel the p value for $\beta = \text{BINOMDIST}(x, n, p_d + (1-p_d)*(1/2), \text{TRUE})$ for x correct responses from n panellists and p_d maximum allowable proportion of discriminators expressed as decimal, i.e. 10% = 0.10. Note that for similarity testing you accept the null hypothesis of no difference with 100(1 - β)% confidence.

Hence, if the p value (for β) is equal to 0.05, you can conclude that the two samples are similar with 95% confidence.

As this Standard is reproduced from an international standard, the following applies:

- (i) Its number appears on the cover and title page while the international standard number appears only on the cover
- (ii) In the source text ‘this International Standard’ should read ‘this Australian Standard’.
- (iii) A full point substitutes for a comma when referring to a decimal marker.

References to International standards should be replaced by references to Australian or Australian/New Zealand Standards, as follows:

<i>Reference to International Standard</i>	<i>Australian Standard</i>
ISO	AS
5492 Sensory analysis—Vocabulary	2542 Sensory analysis of foods
	2542.3 Part 3: Glossary of terms
8009 Sensory analysis—General guidance for the design of test rooms	—

The terms ‘normative’ and ‘informative’ have been used in this Standard to define the application of the annex to which they apply. A ‘normative’ annex is an integral part of a Standard, whereas an ‘informative’ annex is only for information and guidance.

CONTENTS

	<i>Page</i>
1	Scope 1
2	Normative references 1
3	Terms and definitions 2
4	Principle 2
5	General test conditions and requirements 3
6	Assessors 3
6.1	Qualification 3
6.2	Number of assessors 4
7	Procedure 4
8	Analysis and interpretation of results 5
8.1	When testing for a difference 5
8.2	When testing for similarity 5
9	Test report 6
10	Precision and bias 6
	Annex A (normative) Tables 7
	Annex B (informative) Examples 12
	Bibliography 19

Currently in preview, click buy full version

AUSTRALIAN STANDARD

Sensory analysis**Method 2.4:
Specific methods—Duo-trio test****1 Scope**

This International Standard describes a procedure for determining whether a perceptible sensory difference or similarity exists between samples of two products. The method is a forced-choice procedure. The method is applicable whether a difference exists in a single sensory attribute or in several attributes.

The method is statistically less efficient than the triangle test (described in ISO 4120) but is easier to perform by the assessors.

The method is applicable even when the nature of the difference is unknown [i.e. it determines neither the size nor the direction of difference between samples, nor is there any indication of the attribute(s) responsible for the difference]. The method is applicable only if the products are fairly homogeneous.

The method is effective for

- a) determining that
 - either a perceptible difference results (duo-trio testing for difference), or
 - a perceptible difference does not result (duo-trio testing for similarity) when, for example, a change is made in ingredients, processing, packaging, handling or storage;
- b) or for selecting, training and monitoring assessors.

Two forms of the method are described:

- the constant-reference technique, used when one product is familiar to the assessors (e.g. a sample from regular production), and
- the balanced-reference technique, used when one product is not more familiar than the other.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 5492:1984, *Sensory analysis — Vocabulary*

ISO 8586:1988, *Sensory analysis — General guidance for the design of test rooms*