

Australian Standard[®]

**The qualitative identification of
surfactant components in
detergents**

This Australian Standard was prepared by Committee CH/11, Methods of Test for Detergents. It was approved on behalf of the Council of Standards Australia on 5 November 1990 and published on 28 March 1991.

The following interests are represented on Committee CH/11:

- Australian Chemical Industry Council
 - Australian Consumers Association
 - Chemical Confederation of Australia
 - CSIRO, Division of Food Processing
 - CSIRO, Division of Wool Technology
 - Department of Primary Industries and Energy (Commonwealth)
 - The Royal Australian Chemical Institute
 - The University of New South Wales
-

Review of Australian Standards. To keep abreast of progress in industry, Australian Standards are subject to periodic review and are kept up to date by the issue of amendments or new editions as necessary. It is important, therefore, that Standards users ensure that they are in possession of the latest edition, and any amendments thereto.

Full details of all Australian Standards and related publications will be found in the Standards Australia Catalogue of Publications; this information is supplemented each month by the magazine 'The Australian Standard', which subscribing members receive, and which gives details of new publications, new editions and amendments, and of withdrawn Standards.

Suggestions for improvements to Australian Standards, addressed to the head office of Standards Australia, are welcomed. Notification of any inaccuracy or ambiguity found in an Australian Standard should be made without delay in order that the matter may be investigated and appropriate action taken.

This Standard was issued in draft form for comment as DR 81284.

Australian Standard[®]

**The qualitative identification of
surfactant components in
detergents**

First published as AS 3867—1991.

PUBLISHED BY STANDARDS AUSTRALIA
(STANDARDS ASSOCIATION OF AUSTRALIA)
1 THE CRESCENT, HOMEBUSH, NSW 2140

ISBN 0 7262 6647 7

PREFACE

This Standard was prepared by the Standards Australia Committee on Methods of Test for Detergents under the supervision of the Chemical Standards Board.

It was originally requested by the detergents industry because of the growth in scale and complexity of detergents in the market place and to provide a uniform basis for identifying components by separating them from one another systematically. The scheme thus provided is seen as the logical first step in the complete analysis of detergents.

CONTENTS

	<i>Page</i>
1 SCOPE	3
2 APPLICATION	3
3 REFERENCED DOCUMENTS	3
4 DISCUSSION OF SCHEME OF ANALYSIS	3
5 PRINCIPLE	3
 APPENDICES	
A BIBLIOGRAPHICAL REFEREN ES	5
B INFRA-RED SPECTRAL CHARACTERISTICS OF SURFACTANT CONSTITUENTS .	6
C RECOMMENDED METHOD OF ISOLATION OF ACID HYDROLYSIS PRODUCTS	7

© Copyright — STANDARDS AUSTRALIA

Users of Standards are reminded that copyright subsists in all Standards Australia publications and software. Except where the Copyright Act allows and except where provided for below no publications or software produced by Standards Australia may be reproduced, stored in a retrieval system in any form or transmitted by any means without prior permission in writing from Standards Australia. Permission may be conditional on an appropriate royalty payment. Requests for permission and information on commercial software royalties should be directed to the head office of Standards Australia.

Standards Australia will permit up to 10 percent of the technical content pages of a Standard to be copied for use exclusively in-house by purchasers of the Standard without payment of a royalty or advice to Standards Australia.

Standards Australia will also permit the inclusion of its copyright material in computer software programs for no royalty payment provided such programs are used exclusively in-house by the creators of the programs.

Care should be taken to ensure that material used is from the current edition of the Standard and that it is updated whenever the Standard is amended or revised. The number and date of the Standard should therefore be clearly identified.

The use of material in print form or in computer software programs to be used commercially, with or without payment, or in commercial contracts is subject to the payment of a royalty. This policy may be varied by Standards Australia at any time.

STANDARDS AUSTRALIA

Australian Standard

The qualitative identification of surfactant components in detergents

1 SCOPE This Standard sets out a scheme of analysis for the qualitative identification of surfactant components in commercial detergent compositions and in household and personal care products used in Australia. The biodegradability or otherwise of detergents may also be estimated using this scheme. The scheme may also have limited application for other purposes.

NOTE: A list of references for additional reading is included as Appendix A. The use of these references, as appropriate, is recommended for supplementary information.

2 APPLICATION This Standard is for use by laboratories that are required to characterize detergents for comparison, assessment of type or biodegradability characteristics, and which possess facilities for infra-red spectroscopic analysis.

NOTE: This Standard considers only those surfactant components commonly found in detergents in Australia.

3 REFERENCED DOCUMENTS The following documents are referred to in this Standard:

AS

2162 Code of practice for the use of volumetric glassware

2164 One-mark volumetric flasks

4 DISCUSSION OF SCHEME OF ANALYSIS

4.1 General The scheme of qualitative analysis is illustrated by the flow diagram given in Figure 1. This shows the separation of the constituents of the composition into their major component groupings, using solubility and ion characteristics, and the subsequent characterization of individual components, using spectrographic and chromatographic techniques. Guidelines are here given for infra-red spectroscopy because of its wide availability.

NOTE: Guidance is given in Appendix B on interpreting the infra-red absorption characteristics of anionic and cationic constituents of detergents. Typical absorption band characteristics are shown for these components.

4.2 Constituents for identification There are four categories of constituent which may present themselves for identification in detergents. These are described as follows, under their generic headings.

- (a) *Anionic constituents* Anionic constituents are either hydrolysable to acid or non-hydrolysable to acid. Constituents included are as follows:
- Acid hydrolysable anionic constituents. These are alkyl sulfate or alkyl ethoxy sulfate, or both, or alkyl phenol ethoxy sulfate.
 - Acid non-hydrolysable anionic constituents. These are alkyl sulfonate or alkylbenzene sulfonate, or both.

NOTES:

- A method for the isolation and identification of acid-hydrolysable anionic constituents is specified in Appendix C.
- Although soap, which may be present, is an anionic constituent, it is not included above as it appears in a separate fraction during the analysis.

- (b) *Non-ionic constituents* Non-ionic constituents most usually included consist of the following types:

- Alcohol ethoxylate(s), alkylphenol ethoxylate(s), fatty acid ethoxylate(s).
- Fatty acid alkanolamide, fatty acid alkanolamide ethoxylate(s).
- Cyclitol(s), glycerol.
- Polyoxyethylene glycol (PEG), polyoxypropylene glycol (PPG), copolymers containing both polyoxyethylene and polyoxypropylene derivatives.
- Fatty alcohol, fatty alcohol ethoxylate from the corresponding sulfate anionics, unsulfonated alkane, alkylbenzene.

- (c) *Cationic constituents* The common cationic constituents are:

- Alkanolamines; present either as an added component or present as neutralizing agents for the anionic constituent.
- Quaternary ammonium compounds.

- (d) *Amphoteric constituents* These constituents include betaines and amine oxides.

NOTE: Amphoteric constituents tend to be eluted together with the cationic constituents.

5 PRINCIPLE The detergent material is separated into its major component groupings, in accordance with the flow diagram given in Figure 1. The separate groupings are then characterized by infra-red spectroscopy.