



BSI Standards Publication

**Ambient air — Report on  
nitro- and oxy-PAHs — Origin,  
toxicity, concentrations and  
measurement methods**

**National foreword**

This Published Document is the UK implementation of CEN/TR 16998:2016.

The UK participation in its preparation was entrusted by Technical Committee EH/2, Air quality, to Subcommittee EH/2/3, Ambient atmospheres.

A list of organizations represented on this committee can be obtained on request to its secretary.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

© The British Standards Institution 2016.  
Published by BSI Standards Limited 2016

ISBN 978 0 580 91098 2  
ICS 13.040.20

**Compliance with a British Standard cannot confer immunity from legal obligations.**

This Published Document was published under the authority of the Standards Policy and Strategy Committee on 31 December 2016.

**Amendments/corrigenda issued since publication**

Date	Text affected
------	---------------

---

TECHNICAL REPORT

**CEN/TR 16998**

RAPPORT TECHNIQUE

TECHNISCHER BERICHT

November 2016

ICS 13.040.20

English Version

## Ambient air - Report on nitro- and oxy-PAHs - Origin, toxicity, concentrations and measurement methods

Air ambiant - Rapport sur les nitro- et oxy-HAP -  
Origine, toxicité, concentrations et méthodes de  
mesure

Außenluft - Bericht über Nitro- und oxy-PAHs -  
Herkunft, Toxizität, Konzentrationen und  
Messverfahren

This Technical Report was approved by CEN on 21 October 2016. It has been drawn up by the Technical Committee CEN/TC 264.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION  
COMITÉ EUROPÉEN DE NORMALISATION  
EUROPÄISCHES KOMITEE FÜR NORMUNG

**CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels**

<b>Contents</b>		Page
European foreword.....		3
Introduction .....		4
1	Scope.....	5
2	Symbols and abbreviations .....	5
3	Literature overview .....	5
4	Conclusions .....	20
5	Recommendations .....	20
Annex A (informative) Sampling and analysis by GC-MS of some nitro- and oxy-PAHs associated to ambient particulate matter .....		22
A.1	Sampling.....	22
A.2	Analytical materials .....	22
A.2.1	Glassware and sample handling .....	22
A.2.2	Reagents and Solvents.....	22
A.2.3	Extraction apparatus and materials.....	23
A.2.4	Evaporation apparatus and materials.....	24
A.2.5	Clean-up Material.....	24
A.2.6	Weighting Apparatus.....	24
A.2.7	Analytical system .....	24
A.3	Extraction.....	24
A.4	Clean-up.....	24
A.5	Analysis.....	25
A.6	Results.....	25
A.7	Quality assurance .....	30
Annex B (informative) Carcinogenicity and references to nitro- and oxy-PAHs.....		31
Annex C (informative) Mutagenicity of nitro-PAHs .....		34
Annex D (informative) Diesel exhaust data.....		35
Annex E (informative) Structures of nitro- and oxy-PAHs referred in this Technical Report .....		36
Bibliography.....		43

## European foreword

This document (CEN/TR 16998:2016) has been prepared by Technical Committee CEN/TC 264 "Air quality", the secretariat of which is held by DIN.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

Currently in preview, click buy full version

## Introduction

Nitro-PAHs and oxy-PAHs are found in ambient air samples and there are strong indications that they are as harmful as PAHs. Several compounds are classified as probably carcinogenic for humans (see Table in Annex A) and nitro-PAHs are reported to be strongly mutagenic. Photooxidation of volatile PAHs gives rise to the formation of secondary aerosols (Chan et al. 2009, Kautzman et al. 2010, Shakya and Griffin, 2010).

1-Nitropyrene and 2-nitrofluorene are discussed as marker compounds for diesel exhaust and other combustion processes. 2-Nitropyrene and 2-nitrofluoranthene are good marker substances for the formation of nitro-PAHs by secondary reactions.

This Technical Report presents the state of the art of the oxy- and nitro-PAHs topics.

## 1 Scope

This Technical Report is focused on the presence of nitro- and oxy-PAHs in ambient air. It describes how nitro- and oxy-PAH are formed, what typical concentrations are found, what is known about their toxicity, and what sampling and measurement techniques are available.

The conclusions of this report are that nitro- and oxy-PAHs concentrations are present in the atmosphere in levels that are of concern regarding their high toxicity. Information on the presence of these compounds in ambient air is as relevant as information about PAHs. Validated techniques for the measurement of nitro- and oxy-PAHs are available.

## 2 Symbols and abbreviations

DNA	Deoxyribonucleic acid
EI	Electron ionization
CD	Chemiluminescence detection
FD	Fluorescence detection
GC-MS	Gas chromatography – mass spectrometry
GC-NICI-MS	Gas chromatography – negative ion chemical ionization – mass spectrometry
HPLC	High performance liquid chromatography
HPLC-FD	HPLC – fluorescence detection
HPLC-CD	HPLC – chemiluminescence detection
IARC	International Agency for Research on Cancer
LC	Liquid chromatography
MS	Mass spectrometry
NICI	Negative ion chemical ionization
Nitro-PAHs	Nitrated polycyclic aromatic hydrocarbons
Oxy-PAHs	Oxygenated polycyclic aromatic hydrocarbons
PAHs	Polycyclic aromatic hydrocarbons
SPE	Solid phase extraction
ToF-MS	Time of flight mass spectrometry

## 3 Literature overview

### 3.1 Nitro-PAHs

#### 3.1.1 Sources

##### 3.1.1.1 General

Nitro-PAHs in the atmosphere originate mainly from combustion sources and are produced from both gas and heterogeneous phase reactions of the parent PAHs with atmospheric oxidants such as NO<sub>3</sub>, N<sub>2</sub>O<sub>5</sub>, O<sub>3</sub>, OH and peroxide radicals (Arey et al., 1986; Atkinson et al., 1990; Keyte et al., 2013; Pitts et al., 1985; Pitts Jr et al., 1978) in the presence of nitrogen oxides.