



BSI Standards Publication

Ambient air — Monitoring the effects of genetically modified organisms (GMO) — Pollen monitoring

Part 2: Biological pollen sampling using
bee colonies

National foreword

This Published Document is the UK implementation of CEN/TS 16817-2:2015.

The UK participation in its preparation was entrusted by Technical Committee EH/2, Air quality, to Subcommittee EH/2/3, Ambient atmospheres.

A list of organizations represented on this committee can be obtained on request to its secretary.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

© The British Standards Institution 2015.

Published by BSI Standards Limited 2015

ISBN 978 0 580 87462 8

ICS 07.080; 13.020.99

Compliance with a British Standard cannot confer immunity from legal obligations.

This Published Document was published under the authority of the Standards Policy and Strategy Committee on 1 October 2015.

Amendments/corrigenda issued since publication

| Date | Text affected |
|------|---------------|
|------|---------------|

TECHNICAL SPECIFICATION
SPÉCIFICATION TECHNIQUE
TECHNISCHE SPEZIFIKATION

CEN/TS 16817-2

October 2015

ICS 07.080; 13.020.99

English Version

**Ambient air - Monitoring the effects of genetically
modified organisms (GMO) - Pollen monitoring - Part 2
Biological pollen sampling using bee colonies**

Air ambiant - Surveillance des effets d'organismes
génétiquement modifiés (OGM) - Surveillance du
pollen - Partie 2 : Échantillonnage biologique du pollen
à l'aide de colonies d'abeilles

Außenluft - Monitoring der Wirkungen von
gentechnisch veränderten Organismen (GVO) -
Pollenmonitoring - Teil 2: Biologische Pollensammlung
mit Bienenstöcken

This Technical Specification (CEN/TS) was approved by CEN on 16 May 2015 for provisional application.

The period of validity of this CEN/TS is limited initially to three years. After two years the members of CEN will be requested to submit their comments, particularly on the question whether the CEN/TS can be converted into a European Standard.

CEN members are required to announce the existence of this CEN/TS in the same way as for an EN and to make the CEN/TS available promptly at national level in an appropriate form. It is permissible to keep conflicting national standards in force (in parallel to the CEN/TS) until the final decision about the possible conversion of the CEN/TS into an EN is reached.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

| Contents | | Page |
|------------------------|---|-------------|
| European foreword..... | | 4 |
| Introduction | | 5 |
| 1 | Scope | 6 |
| 2 | Normative references | 6 |
| 3 | Terms and definitions | 6 |
| 4 | Basic principle of the procedure | 8 |
| 5 | Sample matrices | 8 |
| 5.1 | Honey..... | 8 |
| 5.2 | Pollen load | 9 |
| 5.3 | Bee-bread..... | 9 |
| 6 | Sampling procedure..... | 9 |
| 6.1 | General..... | 9 |
| 6.2 | Bee colony and hive..... | 9 |
| 6.3 | Sample site..... | 9 |
| 6.4 | Preparation and assembly..... | 10 |
| 6.5 | Exposure time..... | 10 |
| 6.6 | Sampling dates | 10 |
| 6.7 | Extraction, transport and storage..... | 10 |
| 7 | Palynology | 11 |
| 7.1 | General..... | 11 |
| 7.2 | From sample preparation to embedded slide preparation..... | 11 |
| 7.2.1 | General..... | 11 |
| 7.2.2 | Honey..... | 11 |
| 7.2.3 | Pollen load | 11 |
| 7.2.4 | Bee bread | 11 |
| 7.3 | Microscopic analysis..... | 12 |
| 7.4 | Pollen diversity | 12 |
| 8 | Molecular-biological analysis | 13 |
| 8.1 | General..... | 13 |
| 8.2 | Sample preparation..... | 13 |
| 8.2.1 | Honey..... | 13 |
| 8.2.2 | Pollen loads..... | 13 |
| 8.2.3 | Bee bread | 14 |
| 9 | Determination of the target parameters for GMO monitoring and representation of the results | 14 |
| 9.1 | Microscopic pollen analysis | 14 |
| 9.1.1 | General..... | 14 |
| 9.1.2 | Concentration in counts per gram of sample mass..... | 14 |
| 9.1.3 | Total number of pollen collected per exposure time and season..... | 15 |
| 9.1.4 | Relative frequency..... | 15 |
| 9.2 | Molecular-biological analysis | 16 |
| 10 | Performance characteristics of the methods..... | 16 |
| 10.1 | General..... | 16 |

| | | |
|--|---|----|
| 10.2 | Bee colony | 16 |
| 10.3 | Foraging distance | 17 |
| 10.4 | Honey samples, bee-bread samples and pollen load samples | 17 |
| 10.5 | Microscopic pollen analysis | 17 |
| 10.6 | Molecular-biological analysis..... | 17 |
| 11 | Quality assurance and quality control | 18 |
| 11.1 | General measurement strategy and task of pollen monitoring with biological samplers..... | 18 |
| 11.2 | Site protocol..... | 18 |
| 11.3 | Accompanying documentation for samples..... | 19 |
| 11.4 | Parallel measurements..... | 19 |
| 11.5 | Quality assurance and reference materials | 19 |
| Annex A (normative) Maize-specific requirements | | 20 |
| A.1 | Scope | 20 |
| A.2 | Basic principles | 20 |
| A.3 | Sampling | 21 |
| A.4 | Sample preparation | 22 |
| A.5 | Molecular-biological analysis of maize DNA using PCR..... | 23 |
| A.5.1 | General | 23 |
| A.5.2 | DNA extraction..... | 23 |
| A.5.3 | Real-time PCR analysis | 24 |
| A.6 | Determination of the target parameters for GMO monitoring and assessment of the results | 24 |
| Annex B (normative) Rapeseed specific requirements | | 25 |
| B.1 | Scope | 25 |
| B.2 | Basic principles | 25 |
| B.3 | Sampling | 26 |
| B.4 | Sample preparation | 27 |
| B.5 | Molecular-biological analysis of rapeseed DNA using PCR for GMO detection | 28 |
| B.5.1 | General | 28 |
| B.5.2 | DNA extraction..... | 28 |
| B.5.3 | Real-time PCR analysis | 29 |
| B.6 | Determination of the target parameters for GMO monitoring and assessment of the results | 29 |
| Annex C (informative) Good beekeeping practice | | 30 |
| Bibliography | | 31 |

European foreword

This document (CEN/TS 16817-2:2015) has been prepared by Technical Committee CEN/TC 264 "Air quality", the secretariat of which is held by DIN.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

CEN/TS 16817, *Ambient air — Monitoring the effects of genetically modified organisms (GMO) — Pollen monitoring*, is composed of the following parts:

- *Part 1: Technical pollen sampling using pollen mass filter (PMF) and Sigma-2-sampler;*
- *Part 2: Biological pollen sampling using bee colonies* [the present document].

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to announce this Technical Specification: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Introduction

The European Parliament and the European Council require an environmental risk assessment and a post-marketing monitoring for any GMO released to the environment [6; 7]. This had to be implied in national law in any member state of the EC by date.

Pollen dispersal plays a significant role in the dissemination of genetically modified organisms (GMO). Hence, a monitoring procedure that involves recording and documentation of input and distribution of GMO via pollen in a monitoring network mirroring the natural environment is required. For this, technical (CEN/TS 16817-1) and biological sampling of pollen as well as PCR-screening (polymerase chain reaction) procedures are employed to provide evidence of GMO-exposure. The biological sampling system using honey bee colonies is described in the present Technical Specification.

VDI/Guideline 4330 Part 1 [3] presents the necessary fundamentals for the understanding of this Technical Specification. The sampling of pollen in the sample matrices honey, pollen load and bee-bread [5] needs to be viewed in conjunction with the technical sampling for the GMO-monitoring [4].

The use of the biological, actively foraging honeybee and the technical passive samplers complement each other in a manifold and positive way for pollen monitoring of GMO. Therefore it is reasonable to use both. The technical sampling (CEN/TS 16817-1) is based on stationary point-samplers [1]. They give a record of pollen exposure in the air at the sample site that correlates with the prevailing wind direction and relative position to the surrounding pollen sources. The biological sampling using honey bee colonies serves as indicator for GMO exposure in an area and for exposure to roaming insects. Bees display a spatially averaging sampling activity, which represents a cross section of the established, blossoming plants in the area according to the bees collection activities. A wide spectrum of pollen species is recorded using both sampling methods with the procedures complementing each other across the vegetation period [21].

1 Scope

This Technical Specification describes a procedure through which pollen – in particular pollen of genetically modified organisms (GMO) – can be sampled by means of bee colonies.

Bee colonies, especially the foraging bees, actively roam an area and are therefore area related samplers. Pollen sampling depends on the collection activity of the bees and the availability of pollen sources within the spatial zone according to the bees' preferences (supply of melliferous plants). A colony of bees normally forages over an area of up to 5 km radius (median 1,6 km, mean 2,2 km), in rare cases some bees may also forage in greater distances up to 10 km and more [26].

Foragers fix the gathered pollen on the outside of their hind legs (pollen loads, also known as pollen pellets). Inside the hive they place these pollen loads into comb cells close to the brood nest (bee bread). Furthermore, foragers gather nectar and honeydew. Nectar contains pollen which fell from the anthers of the blossom into the nectar drop, or pollen which was dispersed by the wind and stuck in the nectar of other blossoms or adheres to the sticky honeydew of plants. Nectar and honeydew are converted to honey and stored by the bees in the beehive.

Honey, pollen load and bee-bread may be used as sample matrices for the subsequent analysis of pollen as it is possible to concentrate sufficient amounts of pollen for microscopic and molecular biological diagnostics.

Microscopic analysis is used to identify the various pollen types and to quantify the exposure to the target pollen types in question. GMO exposure is analysed by molecular-biological methods: For analysis of pollen DNA quantitative PCR methods are used and described here in this Technical Specification. The analysis of GMO specific proteins and toxins in pollen is possible, too, using ELISA, but to this date the method has not been evaluated enough in pollen matrices for standardization in this Technical Specification.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the reference document (including any amendments) applies.

DIN 10760, *Analysis of honey — Determination of the relative frequency of pollen*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

bee bread

pollen load stored in comb cells close to the brood nest

3.2

bee colony

colony of the honeybee species *Apis mellifera*

3.3

hive

container in which honeybees are kept by beekeepers