



BSI Standards Publication

**Nanotechnologies — Electron spin resonance (ESR)
as a method for measuring reactive oxygen species
(ROS) generated by metal oxide nanomaterials**

National foreword

This Published Document is the UK implementation of ISO/TS 18827:2017.

The UK participation in its preparation was entrusted to Technical Committee NTI/1, Nanotechnologies.

A list of organizations represented on this committee can be obtained on request to its secretary.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

© The British Standards Institution 2017
Published by BSI Standards Limited 2017

ISBN 978 0 580 83155 3

ICS 07.120

Compliance with a British Standard cannot confer immunity from legal obligations.

This Published Document was published under the authority of the Standards Policy and Strategy Committee on 31 July 2017.

Amendments/corrigenda issued since publication

Date	Text affected
------	---------------

**Nanotechnologies — Electron spin
resonance (ESR) as a method for
measuring reactive oxygen species
(ROS) generated by metal oxide
nanomaterials**

*Nanotechnologies — Résonance paramagnétique électronique (RPE)
pour la mesure des espèces réactives de l'oxygène (ROS) générées par
des nanomatériaux sous forme d'oxyde métallique*





COPYRIGHT PROTECTED DOCUMENT

© ISO 2017. Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Ch. de Blandonnet 8 • CP 401
CH-1214 Vernier, Geneva, Switzerland
Tel. +41 22 749 01 11
Fax +41 22 749 09 47
copyright@iso.org
www.iso.org

Contents

Page

Foreword	v
Introduction	vi
1 Scope	1
2 Normative references	1
3 Terms, definitions and abbreviations	1
3.1 Terms and definitions.....	1
3.2 Abbreviations.....	2
4 Principle	2
4.1 General.....	2
4.2 Spin trapping method.....	2
4.2.1 General.....	2
4.2.2 DMPO.....	2
4.2.3 BMPO.....	3
4.2.4 TPC.....	3
4.3 Positive control for generating free radicals.....	3
4.3.1 Fenton reaction[14].....	3
4.3.2 Hypoxanthine-xanthine oxidase system[15].....	3
4.3.3 Rose bengal photosensitization[16][17].....	4
5 Reagents	4
6 Apparatus	4
7 Sampling	5
7.1 Preparation of test sample (metal oxide nanomaterial suspension).....	5
7.2 Preparation of solution for generating hydroxyl radical.....	5
7.2.1 FeSO ₄ solution.....	5
7.2.2 H ₂ O ₂ solution.....	5
7.3 Preparation of solution for generating the superoxide anion radical.....	5
7.3.1 Phosphate buffer.....	5
7.3.2 Hypoxanthine solution.....	5
7.3.3 Xanthine oxidase solution.....	5
7.4 Preparation of solution for generating the singlet oxygen.....	5
7.5 Preparation of spin trapping agent.....	6
7.5.1 General.....	6
7.5.2 DMPO stock solution.....	6
7.5.3 BMPO stock solution.....	6
7.5.4 TPC stock solution.....	6
7.6 Reaction of test sample and spin trapping agent.....	6
7.6.1 General.....	6
7.6.2 DMPO reaction.....	6
7.6.3 BMPO reaction.....	7
7.6.4 TPC reaction.....	7
7.7 Reaction of positive control and spin trapping agent.....	7
7.7.1 DMPO radical adduct form (DMPO/OH).....	7
7.7.2 BMPO radical adduct form (BMPO/OOH).....	7
7.7.3 TPC radical adduct form (TPC/ ¹ O ₂).....	7
7.8 Preparation of the standard sample for spin calculation.....	7
8 Interferences	8
8.1 Sampling.....	8
8.2 Sampling time.....	8
9 Procedure	8
9.1 General.....	8
9.2 Injection of sample.....	9

9.3	ESR measurement.....	10
10	Examples of results.....	16
10.1	DMPO/OH.....	16
10.2	BMPO/OOH.....	16
10.3	TPC/ 10^2	16
10.4	TEMPOL.....	17
	Bibliography.....	18

Currently in preview, click buy full version

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 229, *Nanotechnologies*.

Introduction

Recently, the use of metal or metal oxide-based nanomaterials has dramatically increased in biomedical and industrial applications. However, the scientific basis for the cytotoxicity and genotoxicity of most manufactured nanomaterials are not fully understood. An important mechanism of nanotoxicity is the generation of reactive oxygen species (ROS). The study on the hazardous effects of metal oxide nanomaterials is still in its initial stage. The ability to generate ROS is one main source of toxicity of metal oxide nanomaterials. Overproduction of ROS can induce oxidative stress, resulting in cells failing to maintain normal physiological redox-regulated functions. This in turn may lead to DNA damage, unregulated cell signalling, change in cell motility, cytotoxicity, apoptosis and cancer initiation. There are critical determinants that can affect the generation of ROS. The critical determinants include size, shape, particle surface, surface positive charges, surface-containing groups, particle dissolution, metal ion release from nanometals and nanometal oxides, UV light activation, aggregation, mode of interaction with cells, inflammation and pH of the medium[1]. Thus, to detect and quantify ROS formation on the surface of metal oxide nanomaterials, this document suggests the electron-spin-resonance (ESR) method.

Amongst ROS, the most biologically relevant and widely studied are hydroxyl radical ($\text{OH}\cdot$), superoxide anion radical ($\text{O}_2\cdot^-$), singlet oxygen ($^1\text{O}_2$) and hydrogen peroxide (H_2O_2).

However, direct detection of some free radicals (e.g. superoxide anion and hydroxyl radical) is very difficult or impossible[2] in solution at room temperature. ESR spin trapping is a valuable tool in the study of transient free radicals[3]. Spin trapping is a technique, developed in the late 1960s, where a nitrene or nitroso compound (a spin trap) reacts with a target free radical to form a stable and distinguishable free radical (spin adducts) to be detected by ESR spectroscopy.

Spin adducts can be observed directly by ESR spectroscopy. The ESR spectra of these spin adducts are unique and provide a fingerprint for the presence of ROS.

This document specifies methods of detection by ESR of 5,5-dimethyl-1-pyrroline-N-oxide (DMPO) hydroxyl adduct, 5-tert-butoxycarbonyl-5-methyl-1-pyrroline-N-oxide (BMPO) superoxide adduct and 2,2,5,5-tetramethyl-3-pyrroline-3-carboxamide (TFC) singlet oxygen adduct formation from metal oxide nanomaterials. This document provides a method to assess ROS generation on the metal oxide nanomaterials in a cell free condition. This method may provide valuable information for the prediction of ROS-mediated cytotoxicity without cytotoxicity assay at physico-chemical evaluation phase.

Nanotechnologies — Electron spin resonance (ESR) as a method for measuring reactive oxygen species (ROS) generated by metal oxide nanomaterials

1 Scope

This document provides a procedure for the detection of ROS (OH , O_2^- , $^1\text{O}_2$) generated by metal oxide nanomaterials in aqueous solution with a reactive oxygen species-specific spin trapping agent using ESR, but excludes ESR procedures that do not use a spin trapping agent.

2 Normative references

There are no normative references in this document.

3 Terms, definitions and abbreviations

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <http://www.iso.org/obp>

3.1 Terms and definitions

3.1.1

nanomaterial

material with any external dimension in the nanoscale or having internal structure or surface structure in the nanoscale

Note 1 to entry: This generic term is inclusive of nano-object and nanostructured material.

Note 2 to entry: See also ISO/TS 80004-1:2015, 2.8 to 2.10.

[SOURCE: ISO/TS 80004-1:2015, 2.4]

3.1.2

test sample

material, device, device portion, component, extract or portion thereof that is subjected to biological or chemical testing or evaluation

[SOURCE: ISO/TS 10993-5:2009, 3.5]

3.1.3

zero baseline control

equivalent of the positive control where no radicals are detected

Note 1 to entry: For example, zero baseline control for the positive control of fenton reaction will be H_2O_2 and DMPO in the absence of iron; for hypoxanthine-xanthine oxidase (HX-XO) system, it will be hypoxanthine and BMPO in the absence of HX-XO; for rose bengal photosensitization, it will be rose bengal and TPC in the absence of light.